arXiv:1202.2262v1 [nucl-th] 10 Feb 2012

"Chemical" composition of the Quark-Gluon Plasma in relativistic heavy-ion collisions

F. Scardina,[1,2] M. Colonna,[1] S. Plumari,[1,3] and V. Greco[1,3]
[1] *INFN-Laboratori Nazionali del Sud, Via S. Sofia 62, I-95123 Catania, Italy*
[2] *Physics Dept., University of Messina, Italy*
[3] *Physics and Astronomy Dept., University of Catania, Italy*

We study the evolution of the quark-gluon composition of the plasma created in ultra-Relativistic Heavy Ion Collisions (uRHIC's) employing a partonic transport theory that includes both elastic and inelastic collisions plus a mean fields dynamics associated to the widely used quasi-particle model. The latter, able to describe lattice QCD thermodynamics, implies a "chemical" equilibrium ratio between quarks and gluons strongly increasing as $T \to T_c$, the phase transition temperature. Accordingly we see in realistic simulations of uRHIC's a rapid evolution from a gluon dominated initial state to a quark dominated plasma close to T_c. The quark to gluon ratio can be modified by about a factor of ~ 20 in the bulk of the system and appears to be large also in the high p_T region.

We discuss how this aspect, often overflown, can be essential for a quantitative study of several key issues in the QGP physics: shear viscosity, jet quenching, quarkonia suppression. Furthemore a bulk plasma made by more than 80% of quarks plus antiquarks provides a theoretical basis for hadronization via quark coalescence.

PACS numbers: 12.38.Mh, 25.75.Nq

The search for the Quark-Gluon Plasma (QGP) started its golden age thanks to the experiments at the Relativistic Heavy Ion Collider (RHIC) that have supplied convincing physical observations that a new state of matter has been created [1, 2]. Such a matter has a very small viscosity [3–5], a high opacity to high$-p_T$ particles [6], a strong screening of the interaction able to significantly dissociate charmonia [7], and exihibits a modification of the hadronization respect to the vacuum toward a quark coalescence mechanism [8–10]. Furthermore some RHIC data hints to the creation of an "exotic" initial state of matter, the Color Glass Condensate, that could be the high-energy limiting state of the QCD interaction [6]. The new and upcoming experiments at the Large Hadron Collider (LHC) have confirmed the main gross properties observed at RHIC [11], but they will allow to explore a larger temperature range also with quite different heavy-quark abundancies and will provide more suitable conditions for creating CGC phase as initial state.

The several probes mentioned above rely on the comparison between experimental data and model predictions. A closer look into the several theorethical approaches to the different QGP probes reveals that in some cases the QGP is described as a Gluon Plasma because initially this should be the case. In fact most of the particles come from low x momentum fraction where the nucleon parton distribution functions are gluon dominated. This for example is the case of the most popular jet quenching models assuming a bulk gluon matter. In other cases as in the viscous hydrodynamics a chemical quark-to-gluon equilibration is implicit in the employment of a lattice QCD (lQCD) equation of state $P(\epsilon)$. For the study of quarkonia instead usually one considers a plasma of gluons or an equilibrated QGP according to a massless quark-gluon description acting for dissociation. On the contrary the observation of quark-number scaling in the elliptic flow and the baryon over meson enhancement are explained by quark coalescence models based on a quark dominance in the plasma [8, 10].

Certainly despite a lack of full integration of the different descriptions of the QGP, all of these have been useful simplifications that allowed to successfully identify the creation of the QGP plasma and its gross properties. Nonetheless once we have identified the main qualitative features of the matter created in uRHIC's a quantitative knowledge of properties like the η/s or the solution to open issues on the jet quenching mechanism, quarkonia dissociation-regeneration, hadronization mechanism and existence of a CGC matter, requires to consider the poorly explored issue of the "chemical" composition of the QGP.

The assumption of chemical equilibrium of the QGP, when considered, is usually discussed treating the QGP as a gas of massless quarks and gluons; therefore the expected ratio is given simply by the ratio of the degrees of freedom $N_{q+\overline{q}}/N_g = d_{q+\overline{q}}/d_g = 2N_c\,N_f/(N_c^2-1) = 9/4$ for a system with 3 flavors. On the other hand, as well known from lQCD, the QGP appears to be significantly different from a mere massless gas, showing deviation of both the energy density and the pressure from the $\epsilon/T^4 = 3P/T^4 = cost.$, and in particular exihibiting a large trace anomaly $\langle\Theta^\mu_\mu\rangle = \epsilon - 3P$. It has been shown that such a behavior can be described in terms of a massive quasiparticle model in which both gluons and quarks acquire a thermal mass $m(T) \sim g(T)\,T$, as suggested also by the Hard Thermal Loop (HTL) approach [12–16] or dimensionally reduced screened perturbation theory (DRSPT) [17, 18].

In particular a quasi-particle model (QPM) with a $T-$dependent Bag constant has been successfully applied to quantitatively describe lQCD results for equilibrium thermodynamics[19–21] including the recent ones performed with an unprecedented level of accuracy at the physical quark masses [22]. We notice that if the QGP can be described in terms of finite mass excitations this has a strong impact on the quasiparticle chemical ratio $N_{q+\overline{q}}/N_g$. In fact at equilibrium one has:

$$\frac{N_{q+\overline{q}}}{N_g} = \frac{d_{q+\overline{q}}}{d_g}\,\frac{m_q^2(T)\,K_2(m_q/T)}{m_g^2(T)\,K_2(m_g/T)}\,, \qquad (1)$$

where K_2 is the Bessel function and $m_{q,g}(T)$ are the $T-$dependent quark and gluon masses that can be determined by a fit [19] to recent lQCD calculations [22]. In Fig.1, we show by solid line the equilibrium ratio when the fit to lQCD $\epsilon(T)$ is done assuming $m_q^2/m_g^2 = 3/2\cdot(N_c^2-1)/N_c(2N_c+N_f) = 4/9$ according to a pQCD scheme [20]. We plot also by dash-dotted line the expected ratio assuming the HTL $m_q^2/m_g^2 = 1/9$ ratio. This of course leads to a larger quark abundancy. Even if in Ref. [19] some of the authors have shown that in the last case one can better describe the diagonal quark susceptibilities, in the following we will consider the more commonly assumed pQCD case, which also prevents from overestimates of the magnitude of the dynamics studied.



FIG. 1: Quark to gluon ratio at equilibrium as a function of the temperature as predicted by QPM [19]. Solid line is assuming a m_q/m_g ratio according to pQCD and dot dashed line according to HTL; by dashed line it is indicated the massless case.

In this Letter we discuss the issue of the quark-to-gluon ratio of the matter created in uRHIC's at both RHIC and LHC energies. We employ a Boltzmann-Vlasov transport theory to simulate the partonic stage of the HIC in a realistic way. In the last years several codes have been developed based on transport theory at the cascade level [23–25, 27], i.e. including only collisions between massless partons, with quite rare exceptions [28, 29]. These approaches have been more recently developed to study the dynamics of the partonic stage of the HIC at fixed shear viscosity [27, 29–31] with the advantages to explore possible effects of kinetic non-equilibrium, having also a wider range of validity in p_T and in η/s.

We present here for the first time results within a transport approach that includes the mean field dynamics associated to the thermal self-energies generating the finite mass $m(T)$ in the quasi-particle model discussed above [19, 20]. The approach is formally similar to the one developed in Ref. [29] for the NJL mean field dynamics, but here the quasiparticle mean field allows to account for the proper equation of state, $P(\epsilon)$, as evaluated from lQCD. In such a picture the relativistic Boltzmann-Vlasov equation can be written as follows:

$$[p^\mu\partial_\mu + m^*(x)\partial_\mu m^*(x)\partial_p^\mu]f(x,p) = \mathcal{C}[f](x,p) \qquad (2)$$

where $\mathcal{C}(x,p)$ is the Boltzmann-like collision integral, main ingredient of the several cascade codes:

$$\mathcal{C} = \int\!\!\!\int\!\!\!\int_{2\,1'2'}(f_{1'}f_{2'} - f_1 f_2)|\mathcal{M}_{1'2'\to 12}|^2\delta^4(p_1+p_2-p_1'-p_2') \qquad (3)$$

where $\int_j = \int_j d^3p_j/(2\pi)^3\,2E_j$, f_j are the particle distribution functions, while $\mathcal{M}_{f\to i}$ denotes the invariant transition matrix for elastic as well as inelastic processes. The elastic processes have been implemented and discussed in several previous works [23–25, 27]. The inelastic processes between quarks and gluons ($gg \leftrightarrow q\overline{q}$) is instead the main focus of the present Letter. We have evaluated the matrix element in a pQCD LO order scheme. The tree diagrams contributing to the $gg \leftrightarrow q\overline{q}$ correspond to the $u,t,s-$channels: $\mathcal{M} = \mathcal{M}_s + \mathcal{M}_t + \mathcal{M}_u$. For the massless case the cross sections for such processes are the textbook pQCD cross section for jet production in high-energy proton-proton collisions. With massive quarks the calculations are the renowed Combridge cross sections [32] used to evaluate heavy quark production. In our case we have considered a finite mass for both gluons and quarks together with a dressed gluon propagator. The details of the calculations are lengthy and will be published elsewhere [34], and are quite similar to the one in [33] for finite current strange quark mass. we only mention that the cross section is dominated by the $t-$ and $u-$channel while the $s-$channel and all the interference terms are neglible. The squared matrix element of the $t-$channel is given by:

$$\begin{aligned}|(t-m_q^2)\,\mathcal{M}_t|^2 \;=\;& \frac{8}{3}\pi^2\alpha_s^2[(m_q^2-t)(m_q^2-u) \\ & -2m_q^2(t+m_q^2) - 4\,m_q^2m_g^2 - m_g^4] \qquad (4)\end{aligned}$$

and of course by crossing symmetry the $u-$channel can be obtained by $u \leftrightarrow t$ exchange.

At the temperature explored by the system created at RHIC and LHC the energy scale of the processes, $Q^2 = (2\pi T)^2$, is of the order of $\sim 4 - 6\,\mathrm{GeV}^2$ therefore the perturbative estimate of the strong coupling is $\alpha_s \sim 0.7 - 0.8$. Nonetheless the pQCD scheme with renormalized fermionic and bosonic lines should provide a useful guideline for the estimate of the pertinent cross section, keeping in mind that it can be expected that the real cross section could be even larger than the estimated ones. This is known to be the case for the elastic scattering processes. However this can only increase the magnitude of the effect that we discuss here. Furthermore the evaluated cross sections are already large enough that their variations within a factor of two only marginally affects the final quark to gluon ratio, in fact the system gets anyway quite close to equilibrium.

The thermodynamical self-consistency of the QPM requires a self-consitency between the Bag constant and the effective mass of the quasiparticles [19] which leads to a gap-like equation coupled to Eq.(2):

$$\frac{\partial B}{\partial m_i} = -\int \frac{d^3\vec{p}}{(2\pi)^3}\frac{m_i(x)}{E_i(x)} f_i(x,p) \tag{5}$$

with $i = q, \overline{q}, g$. Eq.(5) allows to evaluate locally the mass in Eq.(2) also in non-equilibrium conditions guaranteeing the conservation of the energy-momentum tensor of the fluid. For the numerical solutions of Eqs.(2) and (5) we use a three dimensional lattice that discretizes the space and the standard test particle methods that samples the distribution function $f(x,p)$ by means of an ensemble of points in the phase-space, for more details see Ref. [29]. We have carefully checked that the numerical implementation in a stationary box is able to reproduce correctly both the kinetic equilibrium and in particular the chemical one, i.e. the right abundancy of quarks, antiquarks and gluons according to Eq.(1), which implies also a proper implementation of the detailed balance.



FIG. 2: Quark to gluon ratio as a function of time (normalized to τ_0) in $Au+Au$ at $\sqrt{s} = 200$AGeV (black lines) and for $Pb+Pb$ at $\sqrt{s} = 5.5$ATeV (light lines). Dashed lines are for the massless case and the solid for the massive case

We have simulated $Au+Au$ collisions at $\sqrt{s} = 200$ AGeV and $Pb+Pb$ at $\sqrt{s} = 5.5$ATeV for $0-10\%$ centrality. The initial conditions in the r-space are given by the standard Glauber condition while in the p-space we use a Boltzmann-Juttner distribution function up to a transverse momentum $p_T = 2$ GeV while at larger momenta minijet distributions are implemented, as calculated by pQCD at NLO order [8]. At RHIC the initial maximum temperature at the centre of the fireball is $T_0 = 340$ MeV and the initial time $\tau_0 = 0.6$ fm/c (corresponding also to the $\tau_0 \cdot T_0 \sim 1$ criterium) as in succesfull hydrodynamical simulations. For $Pb+Pb$ collisions at $\sqrt{s} = 5.5$ ATeV we have $T_0 = 600$ MeV and $\tau_0 \sim 1/T_0 = 0.3$ fm/c. In Fig.2 it is shown the time evolution of the ratio $R_{qg} = N_{q+\overline{q}}/N_g$ for both the massless case (dashed line) and the massive quasi-particle case (solid line). We clearly see that in the massless case the system reaches very quickly, in less than 1 fm/c, the chemical equilibrium given by $R_{qg} \sim 2$. Therefore in the massless case to assume a chemical equilibrium in modeling the plasma composition can be already considered a reasonable approximation.

When the quasi-particle massive case is considered, we can again see that one can reach quickly a value of $R_{qg} \sim 2.5$, then there is a slower rise that leads to a continue increase up to $R_{qg} \sim 4.5$ for $Au+Au$. We notice that for the massive case the equilibrium value is strongly T dependent, especially close to T_c (see Fig.1), and the system more dilute in this stage is not capable to follow such a rapid change. Nonetheless we find that the fireball reaches a value relatively close to the equilibrium at $T \sim T_c$ and eventually it is composed by about 85% of quark plus anti-quarks. At LHC the trends are very similar but a longer part of the lifetime is spent in a T-region where the equilibrium R_{qg} is nearly constant. This results into a moderately smaller final ratio.

We show in Fig.2 the result starting from an initial gluon dominated plasma with a $R_{qg} = 0.25$ which is already a quite strongly gluon dominated plasma, however changing this initial ratio by a factor of two affects the final ratio by less than a 10%, while the effect we are describing modifies the value of R_{qg} by more than an order of magnitude.

In Fig.3 the p_T dependence of the quark to gluon ratio is shown for the initial distribution (dotted line) and the freeze-out distributions: massless case (dashed line) and massive case (solid line). Black lines are for $Au+Au$ and light ones fore $Pb+Pb$. We see the large difference between the massless and the massive case and also that the net gluon to quark conversion extends up to quite large p_T. The decrease of the ratio with p_T can be expected considering that high$-p_T$ particles can more easily elude the equilibration dynamics. However, in the massive case, we note a quite strong dependence below $p_T = 2$ GeV that has not to be interpreted as a fast detachment from the chemical equilibrium. In fact the p_T-dependence of the ratio can be evaluated analitically at equilibrium and it is given by

$$\frac{dN/d^2p_T|_{q+\overline{q}}}{dN/d^2p_T|_g} = \frac{d_{q+\overline{q}}}{d_g}\frac{m_T^q e^{\gamma[(m_T^q-\beta_0 p_T)/T]}}{m_T^g e^{\gamma[(m_T^g-\beta_0 p_T)/T]}} \tag{6}$$

where β_0 is the radial flow velocity, m_T^q and m_T^g are the transverse masses. We plot in Fig.3 by thin solid line such a function rescaled by a factor 0.85 accounting for the lack of full thermalization. The strong p_T dependence obtained in the transport simulation follows very closely the equilibrium behavior at least up to $p_T \sim 1.5$ GeV. This is a well known effect predicted by hydrodynamics and experimentally observed from SPS to LHC energy for hadronic spectra.



FIG. 3: Quark to gluon ratio vs p_T at the freeze-out time. Labels as in Fig.2. The thin solid line represents the full equilibrium raio, Eq. (6); see text for details.

Our study shows that the most common microscopic description of the lQCD thermodynamics implies a plasma that in uRHIC's evolves from a Glasma toward a quark dominated plasma. The R_{qg} ratio can evolve by more than a factor 20 and at freeze-out is anyway more than a factor 2 larger than the one for an equilibrated massless QGP. Such an effect appears to be quite solid despite uncertainties in the inelastic cross sections and could be even larger and robust if one considers that the quark susceptibilities computed on lQCD hints at even smaller m_q/m_g ratio [19] and hence quite larger equilibrium value of R_{qg}, see dashed-dotted line in Fig.1. Therefore the evolution of quark composition of the QGP should not be discarded for any quantitative studies of the several probes of the QGP and its properties.

Our result provides also a support to the quark coalescence hadronization mechanism, capable to explain two main observations at RHIC: the baryon/meson anomaly and the quark number scaling of the elliptic flow [10]. In fact one of the main criticism to the model was the assumption of a quark plasma, which may appear to be unlikely due to the preconception of a gluon dominated plasma in uRHIC's.

The impact however of the chemical evolution discussed in this Letter appears to be much wider. In fact another key physical issue in the field of the QGP physics is the determination of the shear viscosity. Recently it has been pointed out that a two-component system cannot be treated as an effective one-component system with an average η/s from mixture, but one has to solve hydrodynamics for each component [35]. Therefore the relative abundance of the two components can significantly affect the evaluation of the η/s. Another important issue is related with the particle production at high p_T. Our work shows that a significant gluon to quark conversion happens also at $p_T > 4-5$ GeV in the region where the dominant hadronization mechanism should be independent fragmentation. In this region even if far from chemical equilibrium we still find a modification of R_{qg} of about a factor of 7 respect to modelings discarding the $q \leftrightarrow g$ conversion mechanism. In Fig.4 we show how the ratios, $(p+\overline{p})/2\pi^0$, $(p+\overline{p})/2K$, of identified particle can be modified when the gluon conversion mechanism is included. This kind of observable asks also for a better knoweledge of the fragmentation function (FF) in the relevant region of parton momentum fraction x and Q^2. Here we have used here the AKK FF [36] just to provide a quantitative example of the potential impact of our study. This will be however a subject increasing its relevance with the advent of a new detector for Very High Momentum Particle Identification Detector (VHMPID) for the ALICE experiment at LHC [37].



FIG. 4: Ratio of identified particles from independent fragmentation by AKK parametrizations. The dashed lines are the ratio according to the intial quark and gluon distributions, the solid lines are after the "chemical" evolution.

For high$-p_T$ physics a further important consequence could be the impact on the evaluation of the elastic scattering energy loss and the gluon radiation mechanism responsible for the jet quenching. In fact our study leads to a strong decrease of the gg scattering rates and an increase of the qq and $q\overline{q}$ that however are known to have quite smaller cross sections by a factor $4-8$ depending on the specific channel. Similarly the chemical composition of the QGP affects the physics of quarkonia dissociation in medium. In fact also the scattering rates for $g\, J/\Psi(\Upsilon) \rightarrow c\overline{c}(b\overline{b})$ or $g\, J/\Psi(\Upsilon) \rightarrow gc\overline{c}(b\overline{b})$ processes can be expected to be modified because of the different phase space and the reduced scatterer density. The last aspect has been indeed very recently investigated in Ref.[38] for bottomonium.

In conclusions our study shows that one could expect that the QGP created in uRHIC's, even if it is initially a Glasma should very quickly evolve into a plasma dominated by quark plus antiquarks close to the cross-over temperature T_c. The results are quite robust and devolpments of the QPM [19, 20] or inclusion of three-body inelastic scatterings may even make the effect larger. The result supplies a theorethical justification of the *massive-quark* coalescence hadronization models able to successfully describe several puzzling observations at RHIC and LHC; moreover it can have a wide impact on the main physical issues of the QGP physics as a quantitative estimate of the η/s, the study of the identified high-p_T particles and the related jet quenching mechanism as well as on the physics of the quarkonia suppression. In general quark dominance along with a small η/s and a large opacity to high-p_T minijets shifts the interpretation of the QGP toward even stronger deviations form a perturbative behavior [39] and increased relevance of gluon-radiative energy loss [40].

Acknowledgments

We acknoweledge the support by the MIUR under the FIRB Research Grant and by the ERC under the QGP-Dyn Grant.

[1] Adams J *et al.* (STAR Collaboration), Nucl.Phys. A **757** (2005) 102

[2] Adcox K *et al.* (PHENIX Collaboration), Nucl.Phys. A **757** (2005) 184.

[3] Romatschke P and Romatschke U, Phys.Rev.Lett. **99** (2006) 172301.

[4] H. Song and U. W. Heinz, Phys. Rev. C **77** (2008) 064901

[5] H. Song, S. A. Bass, U. Heinz, T. Hirano and C. Shen, Phys. Rev. Lett. **106** (2011) 192301

[6] M. Gyulassy and L. McLerran, Nucl. Phys. A **750** (2005) 30 [nucl-th/0405013].

[7] R. Rapp, D. Blaschke and P. Crochet, Prog. Part. Nucl. Phys. **65** (2010) 209

[8] Greco V, Ko C and Levai P, Phys. Rev. C **68** (2003) 034904; Greco V, Ko C and Levai P, Phys. Rev. Lett. **90** (2003) 202302.

[9] Fries R, Muller B, Nonaka C and Bass S, Phys.Rev. C **68** (2003) 044902

[10] R. J. Fries, V. Greco, and P. Sorensen, Ann. Rev. Nucl. Part. Sci. **58**, 177 (2008).

[11] J. Schukraft *et al.* (ALICE Collaboration), Journ. Phys. G **38** (2011) 124003 .

[12] E. Braaten and R. D. Pisarski, Phys. Rev. D **45**, R1827 (1992).

[13] J. P. Blaizot and E. Iancu, Nucl. Phys. B **417**, 608 (1994).

[14] J. O. Andersen, E. Braaten, and M. Strickland, Phys. Rev. Lett. **83**, 2139 (1999).

[15] J. P. Blaizot, E. Iancu, and A. Rebhan, Phys. Rev. Lett. **83**, 2906 (1999).

[16] J. O. Andersen, L. E. Leganger, M. Strickland and N. Su, Phys. Lett. B **696** (2011) 468; J. O. Andersen, L. E. Leganger, M. Strickland and N. Su, arXiv:1103.2528 [hep-ph].

[17] K. Kajantie, M. Laine, K. Rummukainen, and Y. Schröder, Phys. Rev. D **67**, 105008 (2003).

[18] J. P. Blaizot, E. Iancu, and A. Rebhan, Phys. Rev. D **68**, 025011 (2003).

[19] S. Plumari, W. M. Alberico, V. Greco and C. Ratti, Phys. Rev. D **84** (2011) 094004

[20] A. Peshier, B. Kampfer, O.P. Pavlenko and G. Soff, Phys. Rev. **D54**, 2399 (1996). P. Levai and U. Heinz, Phys. Rev. **C57**, 1879 (1998) A. Peshier and W. Cassing, Phys. Rev. Lett. **94**, 172301 (2005);M. Bluhm, B. Kämpfer, and G. Soff, Phys. Lett. **B620**, 131 (2005); M. Bluhm and B. Kämpfer, Phys. Rev. **D77**, 114016 (2008).

[21] C. Ratti, R. Bellwied, M. Cristoforetti and M. Barbaro, Phys. Rev. D **85** (2012) 014004

[22] S. Borsanyi *et al.*, JHEP **1011**, 077 (2010)

[23] B. Zhang, C. M. Ko, B. -A. Li and Z. -w. Lin, Phys. Rev. C **61** (2000) 067901

[24] D. Molnar and M. Gyulassy, Nucl. Phys. **A697**, 495 (2002).

[25] Z. Xu and C. Greiner, Phys. Rev. **C71**, 064901 (2005).

[26] Z.-W. Lin *et al.*, Phys. Rev. **C72**, 064901 (2005).

[27] G. Ferini, M. Colonna, M. Di Toro, and V. Greco, Phys. Lett. **B670**, 325 (2009).

[28] W. Cassing and E. L. Bratkovskaya, Nucl. Phys. A **831** (2009) 215.

[29] S. Plumari, V. Baran, M. Di Toro, G. Ferini and V. Greco, Phys. Lett. B **689**, 18 (2010).

[30] V. Greco, M. Colonna, M. Di Toro and G. Ferini, Prog. Part. Nucl. Phys. **62**, 562 (2009).

[31] S. Plumari and V. Greco, arXiv:1110.2383 [hep-ph].

[32] B. L. Combridge, Nucl. Phys. B **151** (1979) 429.

[33] T. S. Biro, P. Levai and B. Muller, Phys. Rev. D **42** (1990) 3078.

[34] F. Scardina, S. Plumari and V. Greco, *in preparation.*

[35] A. El, I. Bouras, F. Lauciello, Z. Xu and C. Greiner, arXiv:1103.4038 [hep-ph].

[36] S. Albino, B.A. Kniehl, G. Kramer, Nucl. Phys. B **803** (2008) 42.

[37] A. Di Mauro, A. Agocs, R. Alfaro, G. G. Barnafoldi, G. Bencze, L. Boldizsar, E. Cuautle and G. De Cataldo *et al.*, Nucl. Instrum. Meth. A **639** (2011) 274.

[38] F. Brezinski and G. Wolschin, Phys. Lett. B **707** (2012) 534

[39] E. Shuryak, Prog. Part. Nucl. Phys. **62** (2009) 48

[40] S. Wicks, W. Horowitz, M. Djordjevic and M. Gyulassy, Nucl. Phys. A **784** (2007) 426